



Emerging infectious disease threats to European herpetofauna

Steven J. R. Allain¹ & Amanda L. J. Duffus²

¹ Durrell Institute of Conservation and Ecology, University of Kent, Canterbury, Kent, UK

² Department of Natural Sciences, School of Nursing, Health, and Natural Sciences, Gordon State College, Barnesville, Georgia, USA

In the past decade, infectious disease threats to European herpetofauna have become better understood. Since the 1990s, three major emerging infections in amphibians have been identified (*Batrachochytrium dendrobatidis*, *B. salamandrivorans*, and ranaviruses) as well as at least one of unknown status (herpesviruses), while two major emerging infections of reptiles (*Ophidiomyces ophiodiicola* and ranaviruses) have been identified in wild European populations. The effects of emerging infections on populations have ranged from non-existent to local extirpation. In this article, we review these major infectious disease threats to European herpetofauna, including descriptions of key mortality and/or morbidity events in Europe of their emergence, and address both the distribution and the host diversity of the agent. Additionally, we direct the reader to newly developed resources that facilitate the study of infectious agents in herpetofauna and again stress the importance of an interdisciplinary approach to examining these infectious diseases.

Keywords: *Batrachochytrium dendrobatidis*, *B. salamandrivorans*, herpesviruses, ophidiomycosis, ranaviruses, amphibians, reptiles

INTRODUCTION

Vertebrates are currently experiencing an ongoing mass extinction event (Ceballos et al., 2017). Infectious diseases are contributing to these declines as they are able to cause major population declines, and can contribute to both local and global extirpations (de Castro & Bolker, 2005). Emerging infections have long been known to pose a serious threat to biodiversity (Scott, 1988; Daszak et al., 2000). The link between the decline and extinction of many amphibian populations with the emergence of infectious agents was drawn approximately 20 years ago (Berger et al., 1998; Daszak et al., 1999). While the link between declines in some reptile populations and emerging infections have been less clear, in the past decade growing evidence shows emerging infectious diseases (e.g. snake fungal disease, Lorch et al., 2016) that are clearly able to decimate populations of these animals as well. Infectious disease related declines can even occur in large, seemingly robust populations of wildlife (Daszak et al., 2003). Therefore, the effects in smaller, more fragile, populations are likely to be even more extreme. Unfortunately, populations of both amphibians and reptiles are shrinking around the globe (Gibbons et al., 2000; Sodhi et al., 2008; Stuart et al., 2004; Todd et al., 2010). This will be exacerbated by other factors such as anthropogenic environmental stressors and infectious disease, which negatively affect wildlife health (Acevedo-Whitehouse & Duffus, 2009).

Duffus and Cunningham (2010) reviewed two major disease threats to European amphibians, *Batrachochytrium dendrobatidis* (*Bd*) and ranaviruses. Since the publication of that review, other emerging infectious diseases in European amphibian populations have been identified, *B. salamandrivorans* (*Bsal*, Martel et al., 2013), and the potentially emerging amphibian herpesviruses (Franklinos et al., 2018; Oraggi et al., 2017). In reptiles, the emergence of two troubling infectious diseases has also been documented in wild populations, snake fungal disease (*Ophidiomyces ophiodiicola*, Franklinos et al., 2017) and ranaviruses (Alves de Matos et al., 2011).

EMERGING INFECTIOUS DISEASE THREATS TO AMPHIBIANS

BATRACHOCHYTRIUM DENDROBATIDIS

The amphibian chytrid fungus (*Batrachochytrium dendrobatidis*, hereafter *Bd*) is a non-hyphal zoosporic chytridiomycete fungus implicated in the decline of amphibian species globally (Skerratt et al., 2007). *Batrachochytrium dendrobatidis* is the causative agent of amphibian chytridiomycosis (Berger et al., 1998) that was first linked to declines in amphibian communities in Central America and Australia. Since then, *Bd* has been found all over the globe and has been associated with the extinction of at least 90 amphibian species, and the declines of several hundred others (Wake & Vredenburg, 2008; Scheele et al., 2019). *Bd* is considered

to be a pandemic pathogen in amphibians (Pasmans et al., 2006) with low genetic diversity among different isolates indicating a rapid spread of the pathogen. Of these strains, the global panzootic lineage (BdGPL) has been attributed to most amphibian declines having likely emerged as a hybrid between less virulent strains (Farrer et al., 2011). For a long time, there was speculation as to where *Bd* had originated, with evidence of America or Africa (Weldon et al., 2004), but recent research shows that the infection originated in east Asia (O’Hanlon et al., 2018).

The standard detection method for *Bd* is to swab an individual with a sterile swab and then test for the presence of *Bd* DNA using real-time PCR (also known as qPCR) after processing the sample (Boyle et al., 2004). Gross clinical signs of *Bd* infection in adult amphibians include, but are not limited to: anorexia, lethargy, unusual skin shedding, reddening of the skin (especially in the vent and upper thigh regions), convulsions and loss of the righting reflex (Fig. 1; Nichols et al., 2011). However, the clinical signs of *Bd* vary among species, their life history stages, and other factors such as environmental conditions. In larval stages, clinical signs of chytridiomycosis are also variable. They include but are not limited to: depigmentation of mouthparts, swimming difficulties, and lethargy (reviewed in van Rooij et al., 2015). Therefore, to properly diagnose chytridiomycosis, histology should be used alongside the standard method of qPCR. The pathogen only infects the keratinized tissues of the body, such as the skin in metamorphosed animals and the mouthparts in larvae (Daszak et al., 1999; Marantelli et al., 2004). There are limited studies on the effects *Bd* has on life history, but Garner et al. (2009a) were able to show that experimental infection decreased the time it took tadpoles (of the common toad, *Bufo bufo*) to reach metamorphosis, but also decreased their weight. At higher doses, there were also higher levels of mortality, but these surprising results show that there may be advantageous effects when tadpoles are infected with *Bd*. For more information on host-fungal interactions of *Bd*, please see Van Rooij et al. (2015), and for further



details on the immune response of amphibians to *Bd*, please refer to Grogan et al. (2018).

Currently the potential impacts of *Bd* on European amphibian communities are not fully understood, but within the past decade multiple studies have started to help illuminate the scientific community (e.g. Bielby et al., 2015; Rosa et al., 2013; Tobler & Schmidt, 2010). The amphibian chytrid fungus has both a wide

Figure 1. A common midwife toad (*A. obstetricans*) that died because of *Bd* caused chytridiomycosis. Note the strange position of the legs, due to muscle spasms and the reddish hue of the vent and thighs. Photo by Frank Pasmans.



Figure 2. The distribution of *Batrachochytrium dendrobatidis* (*Bd*) in wild European amphibian populations.

geographic range and a wide host range within Europe (Table 1 and Fig. 2), with infection first being identified in wild populations in 2001 (Bosch et al., 2001). So far, the susceptibility of species across Europe is not fully understood, but *Bd* has been attributed to declines seen in species including the common midwife toad (*Alytes obstetricans*), common toad (*B. bufo*) and European fire salamander (*Salamandra salamandra*) across the Iberian Peninsula (Bosch et al., 2001; Bosch & Martínez-Solano, 2006). Elsewhere in Europe, infection does not always lead to the development of disease, which may be why we have not seen the mass die-offs of amphibians such as those elsewhere in the world (Berger et al., 1998). Experimentally, some anurans species have been shown to have a limited immunity to *Bd*, whilst others have exhibited tolerance, such as the water frogs (*Pelophylax* spp.), which may be an important factor for host-disease dynamics (Daum et al., 2012; Woodhams et al., 2012).

The apparent lack of mass die-offs within many areas Europe may be attributed to the fact that *Bd* has multiple strains of varying virulence (Fisher et al., 2009). Research in the Netherlands has shown over a seven-year time period, populations of the yellow-bellied toad (*Bombina variegata*) are able to coexist with *Bd* (Spitzen-van der Sluijs et al., 2017). Although for now the toads seem to be safe, this may change in the future due to the effects of climate change and other such environmental effects. For more information on the environmental impact of host-pathogen interactions, which can be hugely deterministic in some cases, see Blaustein et al. (2018) for an in-depth discussion. Experiments have also shown that some salamanders such as those in the genus *Speleomantes* also have some immunity to *Bd* (Pasmans et al., 2013). This combined effect of peptide defence and varying *Bd* strains show that at least for now, some species of European amphibians are protected against the potentially negative effects of *Bd* outbreaks (Fisher et al., 2009).

BATRACHOCHYTRIUM SALAMANDRIVORANS

Table 1. Summary of the countries and amphibian species affected by *Batrachochytrium dendrobatidis* infections in wild European populations.

Location	Common Name	Species	First Report
Albania	Macedonian crested newt	<i>Triturus macedonicus</i>	Vojar et al., 2017
	Yellow-bellied toad	<i>Bombina variegata</i>	Vojar et al., 2017
Austria	Alpine newt	<i>Ichthyosaura alpestris</i>	Sztatecsny & Glaser, 2011
	Smooth newt	<i>Lissotriton vulgaris</i>	Sztatecsny & Glaser, 2011
	European water frog complex	<i>Pelophylax</i> spp.	Sztatecsny & Glaser, 2011
	Fire-bellied toad	<i>B. bombina</i>	Sztatecsny & Glaser, 2011
	Great crested newt	<i>T. cristatus</i>	Sztatecsny & Glaser, 2011
	Italian crested newt	<i>T. carnifex</i>	Sztatecsny & Glaser, 2011
	Yellow-bellied toad	<i>B. variegata</i>	Sztatecsny & Glaser, 2011
Belgium	African clawed frog	<i>Xenopus laevis</i>	Spitzen-van der Sluijs et al., 2014
	American bullfrog	<i>Lithobates catesbeianus</i>	Spitzen-van der Sluijs et al., 2014
	Common midwife toad	<i>Alytes obstetricans</i>	Pasmans et al., 2010
	Common toad	<i>Bufo bufo</i>	Martel et al., 2012
	Marbled newt	<i>T. marmoratus</i>	Spitzen-van der Sluijs et al., 2014
Czech Republic	Alpine newt	<i>Ichthyosaura alpestris</i>	Civiš et al., 2012
	Eurasian toads	<i>Bufo</i> sp.	Baláž et al., 2014
	European water frog complex	<i>Pelophylax</i> spp.	Baláž et al., 2014
	Fire-bellied toad	<i>B. bombina</i>	Civiš et al., 2012
	Great crested newt	<i>T. cristatus</i>	Civiš et al., 2012
	Yellow-bellied toad	<i>B. variegata</i>	Civiš et al., 2012
Denmark	Common frog	<i>Rana temporaria</i>	Scalera et al., 2008
	Edible frog	<i>P. esculentus</i>	Scalera et al., 2008
Finland	Common frog	<i>R. temporaria</i>	Patrelle et al., 2012
France	American bullfrog	<i>L. catesbeianus</i>	Garner et al., 2006
	Common midwife toad	<i>A. obstetricans</i>	Garner et al., 2005
	European water frog complex	<i>Pelophylax</i> spp.	Ouellet et al., 2012
Germany	Agile frog	<i>R. dalmatina</i>	Ohst et al., 2011
	Alpine newt	<i>I. alpestris</i>	Ohst et al., 2011
	Common frog	<i>R. temporaria</i>	Ohst et al., 2011
	Common midwife toad	<i>A. obstetricans</i>	Ohst et al., 2011
	Smooth newt	<i>L. vulgaris</i>	Ohst et al., 2011
	Common toad	<i>B. bufo</i>	Ohst et al., 2011
	European water frog complex	<i>Pelophylax</i> spp.	Ohst et al., 2011
	European spadefoot toad	<i>Pelobates fuscus</i>	Ohst et al., 2011
	European tree frog	<i>Hyla arborea</i>	Ohst et al., 2011
	Fire-bellied toad	<i>B. bombina</i>	Ohst et al., 2011
	Fire salamander	<i>Salamandra salamandra</i>	Ohst et al., 2011
	Great crested newt	<i>T. cristatus</i>	Ohst et al., 2011
	Green toad	<i>Bufo viridis</i>	Ohst et al., 2013
	Moor frog	<i>R. arvalis</i>	Ohst et al., 2011
	Natterjack toad	<i>Epidalea calamita</i>	Ohst et al., 2011
	Palmate newt	<i>L. helveticus</i>	Ohst et al., 2011
Yellow-bellied toad	<i>B. variegata</i>	Ohst et al., 2011	
Greece	Epirus water frog	<i>P. epiroticus</i>	Azmanis et al., 2016
	Green frog	<i>B. viridis</i>	Azmanis et al., 2016
	Marsh frog	<i>P. ridibundus</i>	Azmanis et al., 2016
Hungary	Alpine newt	<i>I. alpestris</i>	Vörös et al., 2018
	Edible frog	<i>P. esculentus</i>	Vörös et al., 2018
	European water frog complex	<i>Pelophylax</i> spp.	Tünde et al., 2012
	Fire-bellied toad	<i>B. bombina</i>	Vörös et al., 2018
	Green toad	<i>B. viridis</i>	Vörös et al., 2018
	Marsh frog	<i>P. ridibundus</i>	Vörös et al., 2018
	Yellow-bellied toad	<i>B. variegata</i>	Tünde et al., 2012
Italy	American bullfrog	<i>Lithobates catesbeianus</i>	Garner et al., 2006
	Apennine toad	<i>B. pachypus</i>	Stagni et al., 2004
	Edible frog	<i>P. esculentus</i>	Adams et al., 2008
	Edible frog complex	<i>Pelophylax</i> spp.	Simoncelli et al., 2005
	Fire-bellied toad	<i>B. variegata</i>	Stagni et al., 2002 (in Bovero et al., 2008)
	Italian agile frog	<i>R. latastei</i>	Garner et al., 2004 (in Bovero et al., 2008)
	Italian alpine newt	<i>I. alpestris apuanus</i>	Zampiglia et al., 2013
	Italian crested newt	<i>T. carnifex</i>	Grasseli et al., 2019
	Italian fire salamander	<i>S. salamandra gigliolii</i>	Zampiglia et al., 2013
	Italian stream frog	<i>R. italica</i>	Zampiglia et al., 2013
	Sardinian brook salamander	<i>Euproctus platycephalus</i>	Bovero et al., 2008
Tyrrhenian painted frog	<i>Discoglossus sardus</i>	Bielby et al., 2009	
Luxembourg	Alpine newt	<i>I. alpestris</i>	Wood et al., 2009
	Edible frog complex	<i>Pelophylax</i> spp.	Wood et al., 2009
	Palmate newt	<i>L. helveticus</i>	Wood et al., 2009
Macedonia	European water frog complex	<i>Pelophylax</i> spp.	Vojar et al., 2017
Montenegro	Common or smooth newt	<i>L. vulgaris</i>	Vojar et al., 2017
	Edible frog complex	<i>Pelophylax</i> spp.	Vojar et al., 2017
	Great crested newt	<i>T. cristatus</i>	González et al., 2019

Location	Common Name	Species	First Report
Netherlands	Alpine newt	<i>I. alpestris</i>	Spitzen-van der Sluijs et al., 2014
	Common midwife toad	<i>A. obstetricans</i>	Spitzen-van der Sluijs et al., 2014
	Common frog	<i>R. temporaria</i>	Spitzen-van der Sluijs et al., 2014
	Common or smooth newt	<i>L. vulgaris</i>	Spitzen-van der Sluijs et al., 2014
	Common toad	<i>B. bufo</i>	Spitzen-van der Sluijs et al., 2014
	Edible frog	<i>P. esculentus</i>	Spitzen-van der Sluijs et al., 2014
	European water frog complex	<i>Pelophylax</i> sp.	Spitzen-van der Sluijs et al., 2014
	European tree frog	<i>Hyla arborea</i>	Spitzen-van der Sluijs et al., 2014
	Pool frog	<i>P. lessonae</i>	Spitzen-van der Sluijs et al., 2014
	Natterjack toad	<i>Epidalea calamita</i>	Spitzen-van der Sluijs et al., 2014
	Yellow-bellied toad	<i>B. variegata</i>	Spitzen-van der Sluijs et al., 2014
Poland	Edible frog	<i>P. esculentus</i>	Kolenda et al., 2017
	European water frog complex	<i>Pelophylax</i> sp.	Sura et al., 2010
	Pool frog	<i>P. lessonae</i>	Kolenda et al., 2017
	Yellow-bellied toad	<i>B. variegata</i>	Kolenda et al., 2017
Portugal	Bosca's newt	<i>L. boscai</i>	Rosa et al., 2017
	Common midwife toad	<i>A. obstetricans</i>	Rosa et al., 2013
	Fire salamander	<i>S. salamandra</i>	Rosa et al., 2017
	Iberian green frog	<i>P. perezi</i>	Rosa et al., 2017
	Iberian tree frog	<i>Hyla molleri</i>	Rosa et al., 2017
	Marbled newt	<i>Triturus marmoratus</i>	Rosa et al., 2017
	Spiny toad	<i>B. spinosus</i>	Rosa et al., 2017
Romania	Common Frog	<i>R. temporaria</i>	Vörös et al., 2013
	Smooth newt	<i>L. vulgaris</i>	Vörös et al., 2013
	Yellow-bellied toad	<i>B. variegata</i>	Vörös et al., 2013
Serbia	Edible frog	<i>P. esculentus</i>	Mali et al., 2017
	Marsh frog	<i>P. ridibundus</i>	Mali et al., 2017
	Pool frog	<i>P. lessonae</i>	Mali et al., 2017
Spain	Betic midwife toad	<i>A. dickhilleni</i>	Bosch et al., 2013
	Bosca's newt	<i>L. boscai</i>	Hidalgo-Vila et al., 2012
	Common midwife toad	<i>A. obstetricans</i>	Bosch et al., 2001
	Common toad	<i>B. bufo</i>	Bosch & Martínez-Solano, 2006
	Fire salamander	<i>S. salamandra</i>	Bosch & Martínez-Solano, 2006
	Iberian green frog	<i>P. perezi</i>	Hidalgo-Vila et al., 2012
	Iberian painted frog	<i>Discoglossus galganoi</i>	Hidalgo-Vila et al., 2012
	Majorcan midwife toad	<i>A. muletensis</i>	Garner et al., 2009a
	Mediterranean tree frog	<i>H. meridionalis</i>	Hidalgo-Vila et al., 2012
	Natterjack toad	<i>E. calamita</i>	Martínez-Solano et al., 2003
	Spanish ribbed newt	<i>Pleurodeles waltl</i>	Hidalgo-Vila et al., 2012
	Southern marbled newt	<i>T. pygmaeus</i>	Hidalgo-Vila et al., 2012
	Western spadefoot toad	<i>Pelobates cultripipes</i>	Hidalgo-Vila et al., 2012
Sweden	Common frog	<i>R. temporaria</i>	Kärverno et al., 2018
	Common toad	<i>B. bufo</i>	Kärverno et al., 2018
	Fire-bellied toad	<i>B. bombina</i>	Kärverno et al., 2018
	Moor frog	<i>R. arvalis</i>	Kärverno et al., 2018
	Natterjack toad	<i>E. calamita</i>	Kärverno et al., 2018
	Variable toad	<i>Bufoles variabilis</i>	Kärverno et al., 2018
Switzerland	Alpine newt	<i>I. alpestris</i>	Tobler et al., 2012
	Common midwife toad	<i>A. obstetricans</i>	Tobler et al., 2012
	European water frog complex	<i>Pelophylax</i> sp.	Tobler et al., 2012
	Palmate newt	<i>L. helveticus</i>	Tobler et al., 2012
UK	African clawed frog	<i>Xenopus laevis</i>	Tinsley et al., 2015
	American bullfrog	<i>Lithobates catesbeianus</i>	Garner et al., 2005
	Natterjack toad	<i>E. calamita</i>	May et al., 2011

Table 2. Summary of the countries and amphibian species affected by *Batrachochytrium salamandrivorans* infections in wild European populations.

Location	Common Name	Species	First Report
Belgium	Alpine newt	<i>Ichthyosaura alpestris</i>	Spitzen-van der Sluijs et al., 2016a
	Fire salamander	<i>Salamandra salamandra</i>	Spitzen-van der Sluijs et al., 2016a
Germany	Fire salamander	<i>S. salamandra</i>	Spitzen-van der Sluijs et al., 2016a
Netherlands	Alpine newt	<i>I. alpestris</i>	Spitzen-van der Sluijs et al., 2016a
	Smooth newt	<i>Lissotriton vulgaris</i>	Spitzen-van der Sluijs et al., 2016a
Spain	Fire salamander	<i>S. salamandra</i>	Martel et al., 2013
	Palmate newt	<i>L. helveticus</i>	González et al., 2019

Table 3. Summary of countries and amphibian species affected by herpesvirus infections in wild European populations.

Location	Common Name	Species	First Report
Germany	Common frog	<i>Rana temporaria</i>	Mutschmann & Scheenweiss, 2008
	Moor frog	<i>R. arvalis</i>	Mutschmann & Scheenweiss, 2008
	European spadefoot toad	<i>Pelobates fuscus</i>	Mutschmann & Scheenweiss, 2008
Italy	Agile frog	<i>R. dalmatina</i>	Bennati et al., 1994
Switzerland	Common frog	<i>R. temporaria</i>	Origgi et al., 2017
	Common toad	<i>Bufo bufo</i>	Origgi et al., 2018
UK	Common frog	<i>R. temporaria</i>	Franklinos et al., 2018

The salamander chytrid fungus (*Batrachochytrium salamandrivorans*, hereafter *Bsal*) was first identified in 2013 after a dramatic loss of European fire salamanders (*Salamandra salamandra*) in the Netherlands (Martel et al., 2013; Spitzen-van der Sluijs et al., 2013). The fungus itself is closely related to *B. dendrobatidis*, having diverged sometime in the late Cretaceous or early Paleogene (Martel et al., 2014). The introduction pathway of the pathogen to Europe is believed to have been via the trade in Asian newt species, with *Bsal* being the predominant chytrid fungus affecting Vietnamese salamanders (Laking et al., 2017). There is also alternative evidence to support the introduction of *Bsal* to Europe through the trade in small-webbed fire-bellied toads (*B. microdeladigitora*) from Vietnam (Nguyen et al., 2017). In lab experiments, Asian newts (*Cynops cyanurus*, *C. pyrrhogaster* & *Paramesotriton deloustali*) were discovered to be potential reservoirs of *Bsal*, and worryingly, most European newts died shortly after infection with *Bsal* (Martel et al., 2014). Recent evidence suggests that one of the factors that makes *Bsal* such a threat to urodelean diversity is the fact that infection does not elicit immunity and anurans, such as the common midwife toad (*A. obstetricans*), may act as intermediary hosts (Stegen et al., 2017). More worryingly, alpine newts (*Ichthyosaura alpestris*) can survive for extended periods of time whilst infected with low doses of *Bsal*, with the ability to even clear infection. This is a concern as alpine newts co-occur with *S. salamandra* whilst *A. obstetricans* does not (Stegen et al., 2017).

The clinical signs of *Bsal* are varied (like those of *Bd*) but are usually seen as anorexia, lethargy, ataxia, and skin lesions (Martel et al., 2013). The skin lesions (Fig. 3) are quite diagnostic and easily identified on adult individuals (although this varies by species); however, they are not always present and tend to occur at the end of pathogenesis when the animal is nearly dead (Martel et al., 2013). This means that detailed histological and genetic analyses are needed to confirm the presence of *Bsal* in a suspected infected individual. Again, the standard technique for detection, like *Bd*, is the use of swabs and qPCR.

In the wild, *Bsal* has been only found in a small number of European countries (Table 2) although the presence of the fungus is known to be more widespread in private collections (Fitzpatrick et al., 2018; Sabino-Pinto et al., 2015). It is vital that private collectors take the necessary

biosecurity protocols when disinfecting wastewater and enclosures. More information can be found in Van Rooij et al. (2017). Despite the potentially devastating effects of *Bsal* on naive populations, *Bsal* has a poor dispersal potential which allows some sub-populations to persist in areas where the pathogen is known to be present (Spitzen-van der Sluijs et al., 2018). As highlighted by Spitzen-van der Sluijs et al. (2018), this provides potential for in-the-field mitigation strategies and also shows the importance of biosecurity to ensure that researchers are not accidentally transferring *Bsal* zoospores between sites.

At this time, *Bsal* is limited to wild populations in




Figure 3. Close up of the skin of a fire salamander (*S. salamandra*) that died because of *Bsal* caused chytridiomycosis. Note the skin erosion with black margins indicated by the white arrows. Photo By: Frank Pasmans.

Germany, Belgium, Spain and the Netherlands (Fig. 4). In *Bsal*'s range the species of focus is the European fire salamander (*S. salamandra*) although other species have also tested positive for *Bsal* (Table 2). In order to track the pathogen's spread through Europe, ongoing surveillance needs to be carried out in areas that may be susceptible to spillover, as well as monitoring vulnerable species. Limited screening in the Czech Republic revealed that *Bsal* was not present in wild or captive salamanders (Baláz et al., 2018), although more samples are needed



Figure 4. The distribution of *Batrachochytrium salamandrivorans* (*Bsal*) in wild European amphibian populations.

to ensure that *Bsal* is not present within the country. Ongoing surveillance is needed to ensure the emergence of *Bsal* is not missed in areas where it has not yet been recorded in the wild. This surveillance can be either active or passive, depending on the time and resources available. Recently in the UK, 2,409 swabs collected from wild newts in 2011 and 43 newts submitted for post-mortem examination (between March 2013–December 2017) all tested negative for *Bsal* (Cunningham et al., 2019). However, there is experimental evidence to suggest that salamanders infected with low loads of *Bsal* may go undetected due to latency in detection via qPCR (Thomas et al., 2018). Due to the nature of the pathogen, screening should be completed regularly, especially of those populations that are deemed most vulnerable to ensure that there are no false negatives. In addition to this, to prevent the spread of the pathogen within Europe from captive collections, we advocate the trade in captive bred, certified disease-free salamanders only. Furthermore, the use of the early warning system developed for *Bsal* in Europe should be used (see below for more information).

HERPESVIRUSES

It is thought that amphibian herpesviruses are widely distributed in Europe (Garner et al., 2013); however, they have received relatively little attention in the literature and therefore we cannot determine its status as emergent, although reports appear to be on the rise. The first disease associated with herpesvirus-like particles in an amphibian species reported in Europe occurred in the early 1990s and was published by Benatti et al. (1994). They reported herpesvirus-like particles in agile frogs (*Rana dalmatina*) from Italy suffering from herpes-like lesions. However, the outbreak was not accompanied by a mortality event (Benatti et al., 1994). Subsequently, amphibian herpesviruses have been reported in Germany,




Figure 5. The distribution of amphibian herpesviruses in wild European amphibian populations.

Switzerland, and the UK in five different species, one of conservation concern (*Pelobates fuscus*, Mutschmann & Scheenweiss, 2008, and see Table 3 and Fig. 5).

The gross clinical signs of amphibian herpesvirus infections appear to be conserved across affected species. Typically, lesions are present on the skin that may be tan (more terrestrial species), grey, or grey-blue in colour (typical of more aquatic species; see Oraggi et al., 2017; Franklinos et al., 2018; Oraggi et al., 2018). These are areas where the skin has become thickened and they may appear on any region of the body (Oraggi et al., 2017; Franklinos et al., 2018; Oraggi et al., 2018). The number of lesions that appear on the skin of affected animals is often variable, ranging from one to covering most of the skin (Oraggi et al., 2017; Franklinos et al., 2018; Oraggi et al., 2018; Fig. 6).

There are two species of amphibian herpesvirus known to be present in Europe and both were first described in Swiss amphibians. Oraggi et al. (2017) identified and characterised *Ranid Herpesvirus 3* (RaHV3) from common frogs (*R. temporaria*) that were suffering from a proliferative dermatitis. Subsequently, Oraggi et al. (2018) described *Bufoid Herpesvirus 1* (BfHV1) in common toads (*B. bufo*) that were also suffering from dermatitis. These are the first amphibian herpesviruses outside of North America to be fully characterised and described in wild caught amphibians. Additionally, Franklinos et al. (2018) found herpesvirus-associated dermatitis in common toads. However, the virus(es) present in UK amphibians was/were not fully characterised, so it is unknown if two different virus species are present.

The threat of amphibian herpesviruses to European amphibian populations is unknown. Their ability to cause infection and subsequent mortality in *P. fuscus* is a cause for concern (Garner et al., 2013). More research is needed to fully understand the effects of viral




Figure 6. (A) Herpesvirus infection in a male common frog (*R. temporaria*) from the UK. Note the grayish nature of the lesions that are distributed on the dorsal aspect of the frog. **(B)** Close up of the herpesvirus lesions on the dorsum of the same male common frog. Note the ‘fluid filled’ appearance of the lesions. Photo by Matthew Chester.

persistence in populations of affected amphibians. It will be important for interdisciplinary teams to investigate amphibian herpesviruses because of the difficulties that are often encountered with proper identification and diagnosis (see Garner et al., 2013 for further discussion).

RANAVIRUSES

Amphibian ranaviruses are widely distributed in Europe. They are currently known to be present in over ten countries and in over ten amphibian species (see Table 4 and Fig. 7). Amphibian ranaviruses belong to the viral family *Iridoviridae*, which are large, double stranded DNA viruses (Chinchar et al., 2017). In fact, the first documented cases of what were likely to be *Ranavirus* infections were broadly identified as iridovirus-like agents. Fijan et al. (1991) found iridovirus-like particles in edible frogs (*Pelophylax esculentus*) from a population that experienced a mortality event in Croatia. This discovery was quickly followed by the identification of an iridovirus-like agent associated with morbidity and mortality events in UK common frogs (*Rana temporaria*), that was eventually classified as a *Ranavirus* (Cunningham et al., 1993; Cunningham et al., 1996; Drury et al., 1995). Despite the early report of an iridovirus-like agent in Croatia, no subsequent reports of amphibian ranaviruses can be found in the literature for continental Europe until a new *Ranavirus* species emerged in 2007 (see below for further discussion).

The gross clinical signs of ranaviral disease are conserved across species and life history stages. It is important to note that not all ranaviral infections will result in disease, and therefore the absence of gross clinical signs should not be taken as a clean bill of health (see Rijks et al., 2016). Additionally, in most cases, the gross signs of ranaviriosis are considered to be non-specific to the disease, and therefore it is imperative that a full and proper pathological investigation be performed when infection or disease is thought to be a contributing factor to a morbidity and/or mortality event. Gross clinical signs of ranaviriosis of juvenile and adult amphibians include, but are not limited to: lethargy, skin ulcerations, haemorrhages in the skin, reddening of the skin, necrosis


of digits, and internal haemorrhages of multiple organ systems (predominantly the gastrointestinal tract and reproductive tracts; Cunningham et al., 1996; Cunningham et al., 2007a; Cunningham et al., 2007b; Price et al., 2014; see Figure 8 and images in Duffus & Cunningham, 2010). The signs of disease in affected tadpoles/larvae included systemic external and internal haemorrhages (Balseiro et al., 2009; Balseiro et al., 2010). External haemorrhages occur predominantly in the tissues surrounding the eyes of the tadpoles and in the tail, including the musculature (Balseiro et al. 2009; Balseiro et al., 2010) and edema has also been observed (Rosa et al., 2017). Infection, disease, and mortality can occur at any life history stage in amphibians, except for the egg stage. However, the life history stage affected seems to be species specific. In some species, it is the adults that are predominantly affected [e.g. common frogs (*R. temporaria*, Cunningham et al., 1996)] whereas in other species, the larvae/tadpoles are affected [e.g. common midwife toads (*A. obstetricans*) and Alpine newts (*I. alpestris*) (Balseiro et al., 2009; Balseiro et al., 2010)].

There are several different types of *Ranavirus* that are present in Europe. *Frog virus 3* (FV3), *Common midwife toad virus* (CMTV), and their derivatives appear to be the most common types of ranaviruses present in amphibians. FV3 is the type virus of the genus *Ranavirus* (Tan et al., 2004) and is perhaps the most common species of *Ranavirus* around the globe in amphibian populations (see Duffus et al., 2015). However, CMTV-like viruses are known to recombine with FV3-like strains and can create extremely virulent chimeric viruses (Clayton et al., 2017). Experimental infection of smooth newts (*Lissotriton vulgaris*) with different strains of CMTV *Ranavirus* from the Netherlands clearly demonstrates distinct strain-associated pathogenicity within closely related strains of the virus (Saucedo et al., 2019).

The emergence of amphibian ranaviruses and likely misdiagnosis as ‘red-leg syndrome’ in Europe has had a measurable negative effect on some, but not all, populations. The first well-documented morbidity and mortality event in continental Europe occurred in 2007

Table 4. Summary of countries and amphibian species known to harbour *Ranavirus* infections in wild European populations.

Location	Common Name	Species	First Report
Belgium	Common toad	<i>Bufo bufo</i>	Martel et al., 2012
	American bullfrog	<i>Lithobates catesbeianus</i>	Sharifian-Fard et al., 2011
Croatia	Edible frog	<i>Pelophylax esculentus</i>	Fijan et al., 1991
Denmark	Edible frog	<i>P. esculentus</i>	Ariel et al., 2009
France	Common frog	<i>Rana temporaria</i>	Miaud et al., 2016
Germany	Edible frog	<i>P. esculentus</i>	Stöhr et al., 2013
Italy	Edible frog	<i>P. esculentus</i>	Holopainen et al., 2009
Netherlands	European water frog complex	<i>Pelophylax spp.</i>	Kik et al., 2011
	Smooth newt	<i>Lissotriton vulgaris</i>	Kik et al., 2011
	Common frog	<i>Rana temporaria</i>	Rijks et al., 2016
	Common toad	<i>B. bufo</i>	Rijks et al., 2016
	European spadefoot toad	<i>Pelobates fuscus</i>	Spitzen-van der Sluijs et al., 2016c
Portugal	Great crested newt	<i>Triturus cristatus</i>	Spitzen-van der Sluijs et al., 2016c
	Common midwife toad	<i>Alytes obstetricans</i>	Stöhr et al., 2015
	Bosca's newt	<i>L. boscai</i>	de Matos et al., 2008
	Spiny toad	<i>B. spinosus</i>	Rosa et al., 2017
	Fire salamander	<i>Salamandra salamandra</i>	Rosa et al., 2017
	Iberian tree frog	<i>Hyla molleri</i>	Rosa et al., 2017
Spain	Marbled newt	<i>T. marmoratus</i>	de Matos et al., 2008
	Common midwife toad	<i>A. obstetricans</i>	Balseiro et al., 2009
	Common frog	<i>R. temporaria</i>	Price et al., 2014
	Alpine newt	<i>Ichthyosaura alpestris</i>	Balseiro et al., 2010
	Bosca's newt	<i>Lissotriton boscai</i>	Price et al., 2014
Switzerland	Marbled newt	<i>T. marmoratus</i>	Price et al., 2014
	Marsh frog	<i>P. ridibunda</i>	Stöhr et al., 2015
UK	Common frog	<i>R. temporaria</i>	Drury et al., 1995
	Common toad	<i>B. bufo</i>	Hyatt et al., 2000
	Common midwife toad	<i>A. obstetricans</i>	Duffus et al., 2014
	Smooth newt	<i>L. vulgaris</i>	Duffus et al., 2014

**Figure 7.** The distribution of *Ranavirus* spp. in wild European amphibian populations.

in 'Picos de Europa' National Park, Spain (Balseiro et al., 2009). The mortality event involved common midwife toad (*A. obstetricans*) tadpoles that showed classic signs

of ranavirosis (Balseiro et al., 2009). Investigations into the cause of the mortality event revealed that it was caused by a novel *Ranavirus*, then tentatively called the common midwife toad virus (CMTV, Balseiro et al., 2009) and CMTV was only recently designated as its own species in the genus *Ranavirus* (Chinchar et al., 2017). The following year, CMTV was responsible for another morbidity and mortality event. This time not only did it affect common midwife toad tadpoles, but also alpine newt (*I. alpestris cyreni*) larvae (Balseiro et al., 2010). Since then, declines and local extirpations of several species have been documented in association with the emergence of ranaviruses in the same park. Price et al. (2014) found that at several locations, common midwife toads (*A. obstetricans*) were extirpated and both alpine newts (*I. alpestris*) and common toads (*B. bufo*) experienced severe declines. Unfortunately, it does not appear that the populations are recovering from the effects the emergence of ranaviral infection and disease at this time (Price et al., 2014). Many more outbreaks of ranavirosis have now been documented across continental Europe. Perhaps the largest scale morbidity and mortality event involving a CMTV-like virus occurred in 2010 in the Netherlands (Kik et al., 2011; van Beurden et al., 2014). The event involved more than 1000 individuals, mostly




Figure 8. (A) Ranavirus-associated ulceration on the upper portion of the leg of a common frog (*R. temporaria*) from the UK. (B) Ranavirus-associated ulcerations in a common midwife toad (*A. obstetricans*) from the UK. Arrows indicate the lesions. Note the greyish area around the ulcerations. Photos by Zoological Society of London/Amanda L.J. Duffus



Figure 9. A barred grass snake (*N. helvetica*) in from the UK with suspected ophidiomycosis. (A) Arrow indicates the 'crusty' scales that are common in snakes suffering from ophidiomycosis. (B) Arrows indicate some of the deformed scales frequently seen in snakes with ophidiomycosis. (C) The arrow indicates the crusty scales on the snout. Photos by Silviu Petrovan.


Pelophylax spp. but some common newts (*L. vulgaris*; Kik et al., 2011). Subsequent investigations into amphibian morbidity and mortality events in the Netherlands has revealed that the CMTV-like *Ranavirus* is spreading and is the causative agent of many of the observed events (Rijks et al., 2016; Spitzen-van der Sluijs et al., 2016b; Saucedo et al., 2018). It is also important to note that these outbreaks of disease are affecting the common spadefoot (*P. fuscus*), which is a threatened species (Spitzen-van der Sluijs et al., 2016c). In the UK, common frog populations where ranaviruses have emerged have significantly decreased in size. Teacher et al. (2010) found that on average, common frog populations where *Ranavirus* was found declined by approximately 81 % when compared to populations where the virus was absent. Furthermore, Teacher et al. (2010) report that larger populations were disproportionately affected, with larger populations losing a greater number of animals to disease emergence than those that were initially smaller when *Ranavirus* infections emerged. The emergence of amphibian ranaviruses has also changed the population structure of common frogs in the UK, making them more susceptible to stochastic events (Campbell et al., 2018). This is cause for great concern with the ever-increasing

random events that climate change will cause. Thus, the emergence of ranaviruses in European amphibians is a cause for great conservation concern. They have the potential to be lethal pathogens and are now known to drive populations to decline or even local extinction (e.g. Teacher et al., 2010, Price et al., 2014, Rosa et al., 2017), as a single responsible agent or in co-occurrence with for instance *Bd* and/or *Bsal*. Ranaviruses are persistent pathogens, meaning that they remain on field gear far longer than *Bd* or *Bsal* (Nazir et al., 2012; Van Rooji et al., 2017); therefore, proper disinfection methods should always be used on field gear and enhanced biosecurity in facilities where the pathogen may be found. Investigating amphibian health and disease should be made a priority for many species, even common ones, and include an interdisciplinary approach to ensure that these animals have a future.

The effects of the emergence of ranaviruses may be compounded when another disease, such as *Bd*, has emerged previously in a population. In the 'Serra da Estrela' Natural Park, *Bd* related declines of amphibian populations were occurring for many years before ranavirosis emerged (Rosa et al., 2017). Ranavirosis was first documented in 2011 and is now known to occur in

Table 5. Summary of the countries and snake species affected by snake fungal disease (*Ophidiomyces ophiodiicola*) infections in wild European populations.

Location	Common Name	Species	First Report
Czech Republic	Dice snake	<i>Natrix tessellata</i>	Franklinos et al., 2017
Switzerland	Grass snake	<i>N. natrix</i>	Meier et al., 2018
UK	Barred grass snake	<i>N. helvetica</i>	Franklinos et al., 2017
	Adder	<i>Vipera berus</i>	Franklinos et al., 2017

**Figure 10.** The distribution of ophidiomycosis cases in wild European snake populations.

several species (*T. marmoratus*, *A. obstetricans*, *L. boscai*, *S. salamandra*, and *B. spinus*, Rosa et al., 2017). The emergence of ranaviruses was correlated with sharp declines in two species (*L. boscai* and *A. obstetricans*) at one of the sites within Serra da Estrela Natural Park (Rosa et al., 2017). The ability of *Ranavirus* emergence in amphibians to change the species assemblages at a site and alter host community composition and structure is a huge threat to these animals (Rosa et al., 2017). Therefore, surveillance efforts must not just be centred around one potential pathogen or one potential host; a community-based approach for pathogen surveillance should be used whenever possible.

EMERGING INFECTIOUS DISEASE THREATS TO REPTILES OPHIDIOMYCOSIS/SNAKE FUNGAL DISEASE

In recent years it has become apparent that amphibians are not the only herptiles to suffer from declines caused by emerging fungal pathogens. There are a number of minor fungal diseases that infect reptiles (Paré & Sigler, 2016) but the one that has recently caught international attention is snake fungal disease. The causative agent was first identified in 2009 as *Chrysosporium ophiodiicola* (Rajeev et al., 2009) but subsequent research has seen the fungus moved to its own genus of *Ophidiomyces* (Sigler et al., 2013). Commonly known as snake fungal disease (hereafter SFD) but more correctly known as ophidiomycosis, is troubling for many reasons, one of which is that the clinical signs are not pathognomonic

and so histological evidence, as well as genetic, is needed to confirm infection. SFD was first identified from snakes in the US (Dolinski et al., 2014; Rajeev et al., 2009) and it has recently been found to be present in snake populations in Europe (Franklinos et al., 2017; Meier et al., 2018).

The clinical signs of ophidiomycosis vary slightly between species but include skin lesions, dermatitis, scale deformity and yellow-brown crusty areas (Lorch et al., 2016). Abrasion may create opportunities for infection. Once the stratum corneum has been compromised, *O. ophiodiicola* can then quickly penetrate the epidermis, at which point the host's immune response leads to the aforementioned clinical signs developing (see Fig. 9; Lorch et al., 2015). As a response to the disease, snakes tend to increase their sloughing frequency in order to try to cast off the pathogen. The new skin underneath is clinically normal and this behaviour may lead to snakes successfully curing themselves of ophidiomycosis. However, if old skin is stuck to the new skin during a shed, then there is the potential for reinfection (Lorch et al., 2015). All of this is very worrying for snake species that may already be suffering from the effects of fragmentation, persecution, and inbreeding depression (Madsen et al., 1996; Ursenbacher et al., 2009). The current evidence suggests that *O. ophiodiicola* only has the ability to infect snakes, although this is an area that needs further investigation.

So far, investigations into the presence of ophidiomycosis in Europe are limited, but the presence has been confirmed in a number of species in a small number of countries (Table 5, Fig. 10). The first evidence of ophidiomycosis being present in Europe came from a screening of archived specimens collected by the Garden Wildlife Health project between 2010-2016 (Franklinos et al., 2017). This effort identified that wild European snakes in two countries (the UK and Czech Republic) were positive for *O. ophiodiicola* using a combination of histological and genetic diagnostic techniques. Since then, snakes from Switzerland have also tested positive for the causative agent of ophidiomycosis (Meier et al., 2018). At this stage it is not known how virulent the disease is to European snake species. Franklinos et al. (2017) noted that skin lesions were mild in most of the cases they examined but in some, these lesions were quite severe and would have contributed to the animals' mortality. At this time, we do not have a clear picture of how widespread ophidiomycosis is in Europe or what that means for its snake species. More research is needed to help understand the status of this pathogen. In particular, it is not known whether or not ophidiomycosis is an

Table 6. Summary of countries and reptile species known to harbour *Ranavirus* infections in wild European populations.

Location	Common Name	Species	First Report
Portugal	Iberian mountain lizard	<i>Iberolacerta monticola</i>	Alves de Matos et al., 2011
Spain	Viperine snake	<i>Natrix maura</i>	Price et al., 2014
UK	Slow worm	<i>Anguis fragilis</i>	Price et al., 2017

**Figure 11.** The distribution of *Ranavirus* spp. in wild European reptile species.

endemic disease that has become more virulent due to recent changes or whether it is introduced. Genetic work conducted by Franklino et al. (2017) demonstrates that the strains of ophidiomycosis in Europe are different to those in the US, but again further research is needed to fully understand this relationship.

RANAVIRUSES

Amphibians are not the only group of poikilothermic vertebrates to be affected by ranaviruses; both fish and reptiles are also susceptible. In Europe, most cases of *Ranavirus* infections in reptiles have been reported in captive populations (see Duffus et al., 2015). There are only three reports of *Ranavirus* infections in wild European reptiles in three different species; however, it is extremely likely that this is not reflective of the actual situation (see Table 6 and Fig. 11).

Signs of ranaviruses in reptilians are diverse and non-specific, including but not limited to: lethargy, anorexia, upper respiratory tract necrosis, edema (especially in the region of the neck), dermatitis, and nasal discharge (Marschang, 2011). Most reptilian *Ranavirus* infections have actually been reported in testudinids, with comparatively few in squamates (see Duffus et al., 2015). Since the gross signs of ranaviruses in reptiles are diverse, it is important for a full pathological investigation to be undertaken when disease is thought to be the cause of mortality. The first confirmed report of an iridovirus-like virus was documented in an Iberian mountain lizard (*Lacerta monticola*) from 'Serra da Estrela', Portugal (Alves

de Matos et al., 2011). The animal was asymptomatic and released after a blood sample was taken (Alves de Matos et al., 2011; Stöhr et al., 2015). A *Ranavirus* spp. was isolated from this blood sample and it was shown to be an FV3-like virus (Alves de Matos et al., 2011; Stöhr et al., 2015). A report of a *Ranavirus* infection in a slow worm (*Anguis fragilis*) is made in Price et al. (2017) but very little information about the affected animal was provided. However, in 2014, a more concerning report of a *Ranavirus* infection in a reptile was published. Price et al. (2014) report the first *Ranavirus*-associated mortality in a wild European reptile. A CMTV-like *Ranavirus* was isolated from a dead viperine snake (*Natrix maura*), from 'Picos de Europa' National Park, Spain, that had been scavenging on amphibians from a *Ranavirus*-associated mortality event (Price et al., 2014). This underscores the ability of ranaviruses to infect multiple hosts from different taxa and is a further demonstration of how severe a threat to herpetofauna that this group of pathogens is.

GENERAL RESOURCES FOR STUDYING HERPETOFAUNAL INFECTIOUS DISEASES

In the past decade, there have been a number of important resources published that examine different aspects of the study of herpetofaunal infectious diseases. Gray et al. (2017) provides an in-depth methodology on pathogen surveillance specifically for herpetofauna. They examine how to design a surveillance study, how to collect samples to perform the study, biosecurity, and even intervention strategies for when an emerging infectious disease has been detected. Gray et al. (2017) is a valuable resource for those seeking to improve study design and limit the spread of infectious diseases in herpetofauna. Langwig et al. (2015) provide an excellent resource for examining the emergence of infectious diseases in wildlife. They provide a framework for decision making and necessary actions pre-and post-infection detection.

Few resources for wildlife professionals have been developed that deal with necropsy techniques, as most have been written for veterinary professionals. While we encourage multidisciplinary teams that include veterinary pathologists, necropsies of herpetofauna may also be carried out by trained wildlife professionals. Duffus et al. (2017) provides a guide for investigating herpetofaunal mortality events. The manuscript is geared towards wildlife professionals and provides methods for necropsies, tissue sample collection, and sample storage and shipping. There are also publications that specifically deal with the detection and diagnosis of specific pathogens. For example, Thomas et al. (2018) examines the current diagnostic tools available for *Bsal*. They provide guidelines not only for the use of these

tools, but also for interpreting the results that these tools provide (Thomas et al., 2018). It is incredibly important to standardise both the methodologies and the interpretation of their results to ensure that results from different studies can be easily and consistently compared. Unfortunately, not all studies follow these guidelines and this leads problems in comparing results. For example, Dalbeck et al. (2018) uses a significance threshold of 0.05 GE when testing for *Bsal*, whereas a threshold of 0.1 GE is recommended by Thomas et al. (2018), when diluting extracted samples prior to qPCR. This of course, will lead to confusion and a difference in results when comparing studies unless there is a universally agreed standard, such is recommended in the case of *Bd* (Boyle et al., 2004; Blooi et al., 2013).

In addition to print resources, there are online databases that have been developed to track some emerging infections of herpetofauna. The Amphibian Disease Portal (<https://amphibiandisease.org>) is a database that is used to track reports of *Bd* and *Bsal* around the globe. Additionally, there is a *Bsal* reporting system for Europe (www.BsalEurope.com) and an associated early warning system. The *Bsal* reporting system and the early warning system for Europe will be extremely important in tracking infections and mitigating the effects of *Bsal* once it is found in an area. The Global Ranavirus Reporting System (<https://mantle.io/grrs>) is another online database that tracks reports of *Ranavirus* infection around the globe. These are both relatively new online databases and rely on researchers to upload their data into the system. To ensure that these resources are viable, we encourage researchers to upload their published data to these sites. An older resource that has been previously been used to track the global emergence of *Bd* is www.bd-maps.net (Olson et al., 2013). There are over 40,000 entries from over 80 countries in this database.

Ranaviruses, *Bsal*, and *Bd* are all reportable infections in amphibians (see <http://www.oie.int/animal-health-in-the-world/oie-listed-diseases-2019/> for a list of reportable infections). The OIE (<http://www.oie.int/>) has specific guidelines for reporting these infections and they should be reported to the appropriate agency in the country that they are discovered in. Additionally, risk analyses may be performed and there are several sources for guidelines, including Jakob-Hoff et al. (2014) and one jointly published by the OIE and IUCN (OIE and IUCN 2014). These are invaluable resources for anyone who deals with infectious diseases in wildlife.

CONCLUDING REMARKS

With the global declines of both amphibians and reptiles, it is now more important than ever to understand the threats to these unique and important animals. Although we have only focused on five emerging infectious disease threats to these animals in Europe, there are likely more (e.g. *Amphibiocystidium*, identified in Duffus & Cunningham, 2010), that are understudied and poorly understood, or even unknown infections that are taking a toll on the populations of these animals. Unfortunately, as stated above, the emergence of several of these

infectious diseases has resulted in population declines and local extirpations of several species, some of which were rare, but others, more worryingly, were considered to be common.

An important and necessary difference between the infection status of an animal and the appearance of clinical signs of disease is still often overlooked in many studies in herpetofauna. Infection is the presence of the potentially pathogenic or disease-causing agent in an organism or population, whereas disease is a measurable negative effect that the infection process has imposed in an animal (Scott, 1988). The effects of an infection may or may not result in the development of clinical signs of disease (Scott, 1988). Usually, disease is measured by the appearance of clinical signs. However, the presence/absence of these signs are often dependent upon the scale at which one is examining the organism (Scott, 1988).

Recent research has focused on both the imperfect detectability of infectious diseases (such as *Bd*) when using swabs and qPCR (DiRenzo et al., 2018) as well as the costs involved to complete the analysis (Sabino-Pinto et al., 2018). As previously mentioned, qPCR isn't infallible and false-positive or false-negative results are always possible. We therefore recommend that multiple detection techniques be used in order to be certain when screening suspected infected individuals. In order to fully understand the effects that emerging infectious diseases have on populations, an interdisciplinary team, as stressed in Duffus and Cunningham (2010), should be used. Ideally, such a team would include ecologists, herpetologists, population biologists, and veterinary pathologists. The team should investigate the entire situation, sometimes including multiple species in an area, to garner a deeper understanding of the issues that are potentially being caused by the emergence of an infectious agent.

ACKNOWLEDGEMENTS

We would like to thank Matthew Chester, Silviu Petrovan and Frank Pasmans for the use of their images. We would also like to thank the anonymous reviews, Rachel Antwis and John C. George for comments on an earlier version of the manuscript. Their comments greatly improved the quality of the document.

This work was supported by the Natural Environment Research Council through the EnvEast Doctoral Training Partnership [grant number NE/L002582/1] to S.J.R.A.

REFERENCES

- Acevedo-Whitehouse, K. & Duffus, A. L. J. (2009). Effects of environmental change on wildlife health. *Philosophical Transactions of the Royal Society B* 364, 3429-3438.
- Adams, M. J., Galvan, S., Scalera, R., Grieco, C. & Sindaco, R. (2008). *Batrachochytrium dendrobatidis* in amphibian populations in Italy. *Herpetological Review* 39, 324-326.
- Alves de Matos, A.A., Caeiro, M.F., Marschang, R.E., Papp, T., Soares, C., Marçal, M.R. & Carretero, M.A. (2008). Adaptation of ranaviruses from Peneda-Gerês National Park (Portugal)

- to cell cultures and their characterization. *Microscopy and Microanalysis* 14, 139-140.
- Ariel, E., Kielgast, J., Svart, H.E., Larsen, K., Tapiovaara, H., Jensen, B. B. & Holopainen, R. (2009). *Ranavirus* in wild edible frogs *Pelophylax kl. esculentus* in Denmark. *Diseases of Aquatic Organisms* 85, 7-14.
- Azmanis, P. N., Strachinis, I., Lymberakis, P. & Marschang, R. E. (2016). First detection of the amphibian chytrid fungus (*Batrachochytridium dendrobatidis*) in free-living anuran populations in Greece. *Journal of the Hellenic Veterinary Medical Society* 67, 253-258.
- Baláz, V., Vojar, J., Civi, P. & Rozínek, R. (2014). Chytridiomycosis risk among Central European amphibians based on surveillance data. *Diseases of Aquatic Organisms* 112, 1-8.
- Baláz, V., Solský, M., Lastra González, D., Havlíková, B., Gallego Zamorano, J., González Sevilleja, C., Torrent, L. & Vojar, J. (2018). First survey of pathogenic fungus *Batrachochytrium salamandrivorans* in wild and captive amphibians in the Czech Republic. *Salamandra* 54, 87-91.
- Balseiro, A., Dalton, K. P., del Cerro, A., Marquez, I., Cunningham, A. A., Parra, F., Prieto, J.M. & Casais, R. (2009). Pathology, isolation and characterization of a *Ranavirus* from the common midwife toad, *Alytes obstetricans*, on the Iberian Peninsula. *Diseases of Aquatic Organisms* 8, 950-954.
- Balseiro, A., Dalton, K. P., del Cerro, A., Marquez, I., Parra, F., Prieto, J.M. & Casais, R. (2010). Outbreak of common midwife toad virus in alpine newts (*Mesotriton alpestris cyreni*) and common midwife toad (*Alytes obstetricans*) in northern Spain: a comparative pathological study of an emerging *Ranavirus*. *Veterinary Journal* 186, 256-258.
- Bennati, R., Bonetti, M., Lavazza, A. & Gemetti, D. (1994). Skin lesions associated with herpesvirus-like particles in frogs (*Rana dalmatina*). *Veterinary Record* 135, 625-626.
- Berger, L., Speare, R., Daszak, P., Green, D. E., Cunningham, A. A., Goggin, C. L., Slocombe, R., Ragan, M. A., Hyatt, A. D., McDonald, K. R., & Hines, H. B. (1998). Chytridiomycosis causes amphibian mortality associated with population declines in the rain forests of Australia and Central America. *Proceedings of the National Academy of Sciences* 95, 9031-9036.
- Bielby, J., Bovero, S., Sotgiu, G., Tessa, G., Favelli, M., Angelini, C., Doglio, S., Clare, F. C., Gazzaniga, E., Lapietra, F. & Garner, T. W. J. (2009). Fatal chytridiomycosis in the Tyrrhenian painted frog. *EcoHealth* 6, 27-32.
- Bielby, J., Fisher, M. C., Clare, F. C., Rosa, G.M. & Garner, T. W. J. (2015). Host species vary in infection probability, sub-lethal effects, and costs of immune response when exposed to an amphibian parasite. *Scientific Reports* 5, 10828.
- Blaustein, A., Urbina, J., Snyder, P. W., Reynolds, E., Dang, T., Hoverman, J. T., Han, B., Olson, D. H., Searle, C. & Hambalek, N. (2018). Effects of emerging infectious diseases on amphibians: a review of experimental studies. *Diversity* 10, 81.
- Blooi, M., Pasmans, F., Longcore, J. E., Spitzen-Van Der Sluijs, A., Vercammen, F. & Martel, A. (2013). Duplex real-time PCR for rapid simultaneous detection of *Batrachochytrium dendrobatidis* and *Batrachochytrium salamandrivorans* in amphibian samples. *Journal of Clinical Microbiology* 51, 4173-4177.
- Bosch, J., Martínez-Solano, I. & García-París, M. (2001). Evidence of a chytrid fungus infection involved in the decline of the common midwife toad (*Alytes obstetricans*) in protected areas of central Spain. *Biological Conservation* 97, 331-337.
- Bosch, J. & Martínez-Solano, I. (2006). Chytrid fungus infection related to unusual mortalities of *Salamandra salamandra* and *Bufo bufo* in the Penalara Natural Park, Spain. *Oryx* 40, 84-89.
- Bosch, J., García-Alonso, D., Fernández-Beaskoetxea, S., Fisher, M. C. & Garner, T. W. (2013). Evidence for the introduction of lethal chytridiomycosis affecting wild betic midwife toads (*Alytes dickhilleni*). *EcoHealth* 10, 82-89.
- Bovero, S., Sotgiu, G., Angelini, C., Doglio, S., Gazzaniga, E., Cunningham, A.A. & Garner, T.W.J. (2008). Detection of chytridiomycosis caused by *Batrachochytrium dendrobatidis* in the endangered Sardinian newt (*Euproctus platycephalus*) in southern Sardinia, Italy. *Journal of Wildlife Diseases* 44, 712-715.
- Boyle, D.G., Boyle, D.B., Olsen, V., Morgan, J.A.T. & Hyatt, A.D. (2004). Rapid quantitative detection of chytridiomycosis (*Batrachochytrium dendrobatidis*) in amphibian samples using real-time Taqman PCR assay. *Diseases of Aquatic Organisms* 60, 141-148.
- Campbell, L.J., Garner, T.W.J., Tessa, G., Scheele, B.C., Griffiths, A.G.F., Wilfert, L., & X.A. Harrison (2018). An emerging viral pathogen truncates population age structure in a European amphibian and may reduce population viability. PeerJ 6:e5949 <https://doi.org/10.7717/peerj.5949>.
- Ceballos, G., Ehrlich, P.R., & Dirzo, R. (2017) Biological annihilation via the ongoing sixth mass extinction signaled by vertebrate population losses and declines. *Proceedings of the National Academy of Sciences* 114, E6089-E6096.
- Chinchar, V.G., Hick, P., Ince, I.A., Jancovich, J.K., Marschang, R., Qin, Q., Subramaniam, K., Waltzek, T.B., Whittington, R., Williams, T., Zhang, Q. & ICTV Report Consortium. (2017). *ICTV Virus Taxonomy Profile: Iridoviridae*, *Journal of General Virology* 98, 890-891.
- Civiš, P., Vojar, J., Literák, I. & Baláz, V. (2012). Current state of *Bd* occurrence in the Czech Republic. *Herpetological Review* 43, 75-78.
- Claytor, S.C., Subramaniam, K., Landrau-Giovannetti, N., Chinchar, V.G., Gray, M.J., Miller, D.L., Mavian, C., Salemi, M., Wisely, S., & Waltzek, T.B. (2017). *Ranavirus* phylogenomics: Signatures of recombination and inversions among bullfrog ranaculture isolates. *Virology* 511, 330-343
- Cunningham, A.A., Langton, T.E.S., Bennett, P.M., Drury, S. E. N., Gough, R.E. & Kirkwood, J.K. (1993). Unusual mortality associated with poxvirus-like particles in frogs (*Rana temporaria*). *Veterinary Record* 133, 141-142.
- Cunningham, A.A., Langton, T.E.S., Bennett, P.M., Lewin, J.F., Drury, S.E.N., Gough, R.E. & MacGregor, S.K. (1996). Pathology and microbiological findings from incidents of unusual mortality of the common frog (*Rana temporaria*). *Philosophical Transactions of the Royal Society of London B* 351, 1539-1557.
- Cunningham, A.A., Hyatt, A.D., Russell, P. & Bennett, P.M. (2007a). Emerging epidemic diseases of frogs in Britain are dependent on the source of the *Ranavirus* agent and the route of exposure. *Epidemiology and Infections* 135, 1200-1212.
- Cunningham, A.A., Hyatt, A.D., Russell, P. & Bennett, P.M. (2007b). Experimental transmission of a *Ranavirus* disease of common toads (*Bufo bufo*) to common frogs (*Rana*

- temporaria*). *Epidemiology and Infections* 135, 1213–1216.
- Cunningham, A.A., Smith, F., McKinley, T.J., Perkins, M.W., Fitzpatrick, L.D., Wright, O.N. & Lawson, B. (2019). Apparent absence of *Batrachochytrium salamandrivorans* in wild urodeles in the United Kingdom. *Scientific Reports* 9, 2831.
- Dalbeck, L., Düssel-Siebert, H., Kerres, A., Kirst, K., Koch, A., Lötters, S., Ohlhoff, D., Sabino-Pinto, J., Preißler, K., Schulte U., Schulz, V., Steinfartz, S., Veith, M., Vences, M., Wagner N. & Wegge, J. (2018). Die Salamanderpest und ihr Erreger *Batrachochytrium salamandrivorans* (Bsal): aktueller Stand in Deutschland. *Zeitschrift für Feldherpetologie* 25, 1–22.
- Daszak, P., Berger, L., Cunningham, A.A., Hyatt, A.D., Green, D.E. & Speare, R. (1999). Emerging infectious diseases and amphibian population declines. *Emerging Infectious Diseases* 5, 735–748.
- Daszak, P., Cunningham, A.A. & Hyatt, A.D. (2000) Emerging infectious diseases of wildlife--threats to biodiversity and human health. *Science* 287, 443–449.
- Daszak, P., Cunningham, A.A. & Hyatt, A.D. (2003) Infectious disease and amphibian population declines. *Diversity and Distributions* 9, 141–150.
- Daum, J.M., Davis, L.R., Bigler, L. & Woodhams, D.C. (2012). Hybrid advantage in skin peptide immune defenses of water frogs (*Pelophylax esculentus*) at risk from emerging pathogens. *Infection, Genetics and Evolution* 12, 1854–1864.
- de Castro, F. & Bolker, B. (2005) Mechanisms of disease-induced extinction. *Ecology Letters* 8, 117–126.
- DiRenzo, G.V., Campbell Grant, E.H., Longo, A.V., Che-Castaldo, C., Zamudio, K.R. & Lips, K.R. (2018). Imperfect pathogen detection from non-invasive skin swabs biases disease inference. *Methods in Ecology and Evolution* 9, 380–389.
- Dolinski, A.C., Allender, M.C., Hsiao, V. & Maddox, C.W. (2014). Systemic *Ophidiomyces ophiodiicola* infection in a free-ranging plains garter snake (*Thamnophis radix*). *Journal of Herpetological Medicine and Surgery* 24, 7–10.
- Drury, S.E.N., Gough, R.E. & Cunningham, A.A. (1995). Isolation of an iridovirus-like agent from common frogs (*Rana temporaria*). *Veterinary Record* 137, 72–73.
- Duffus, A.L.J. & Cunningham, A.A. (2010). Major disease threats to European amphibians. *The Herpetological Journal* 20, 117–127.
- Duffus, A.L.J., Fenton, H.M.A., Gray, M.J. & Miller, D.L. (2017). Investigating amphibian and reptile mortalities: A practical guide for wildlife professionals. *Herpetological Review* 48, 550–557.
- Duffus, A.L.J., Garner, T.W.J., Davis, A.R., Dean, A.W. and R.A. Nichols. (2017). Phylogenetic analysis of 24 *Ranavirus* isolates from English amphibians using 2 partial loci. *Journal of Emerging Diseases and Virology* 3, 131.
- Duffus, A.L.J., Nichols, R.A. & Garner, T.W.J. (2014). Detection of a Frog Virus 3-like *Ranavirus* in Native and Introduced Amphibians in the United Kingdom in 2007 and 2008. *Herpetological Review* 45, 608–610.
- Duffus, A.L.J., Waltzek, T.B., Stöhr, A.C., Allender, M.C., Gotesman, M., Whittington, R.J., Hick, P., Hines, M.K. & Marschang, R.E. (2015). Distribution and host range of ranaviruses. In *Ranaviruses*, Gray, M.J., & Chinchar V.G. Eds. Springer.
- Farrer, R.A., Weinert, L.A., Bielby, J., Garner, T.W., Balloux, F., Clare, F., Bosch, J., Cunningham, A.A., Weldon, C., du Preez, L.H. & Anderson, L. (2011). Multiple emergences of genetically diverse amphibian-infecting chytrids include a globalized hypervirulent recombinant lineage. *Proceedings of the National Academy of Sciences* 108(46), 18732–18736.
- Fijan, N., Matašin, Z., Petrinec, Z., Valpotić, I. & Zwillenberg, L.O. (1991). Isolation of an iridovirus-like agent from the green frog (*Rana esculenta* L.). *Veterinarski Arhiv* 61, 151–158.
- Fisher, M.C., Bosch, J., Yin, Z., Stead, D.A., Walker, J., Selway, L., Broun, A.J.P., Walker, L.A., Gow, N.A.R., Stajich, J.E. & Garner, T.W.J. (2009). Proteomic and phenotypic profiling of the amphibian pathogen *Batrachochytrium dendrobatidis* shows that the genotype is linked to virulence. *Molecular Ecology* 18, 415–429.
- Fitzpatrick, L.D., Pasmans, F., Martel, A. & Cunningham, A. A. (2018). Epidemiological tracing of *Batrachochytrium salamandrivorans* identifies widespread infection and associated mortalities in private amphibian collections. *Scientific Reports* 8, 13845.
- Franklinos, L.H.V., Lorch, J.M., Bohuski, E., Fernandez, J.R.R., Wright, O.N., Fitzpatrick, L., Petrovan, S., Durrant, C., Linton, C., Baláž, V. & Cunningham, A.A. (2017). Emerging fungal pathogen *Ophidiomyces ophiodiicola* in wild European snakes. *Scientific Reports* 7, 3844.
- Franklinos, L.H.V., Rodrigues-Ramos Fernandes, J., Hydeskov, H.B., Hopkins, K.P., Everest, D.J., Cunningham, A.A. & Lawson, B. (2018). Herpesvirus skin disease in free-living common frogs *Rana temporaria* in Great Britain. *Diseases of Aquatic Organisms* 129, 239–244.
- Garner, T. W.J., Walker, S., Bosch, J., Hyatt, A.D., Cunningham, A.A. & Fisher, M.C. (2005). Chytrid fungus in Europe. *Emerging Infectious Diseases* 11, 1639–1641.
- Garner, T.W.J., Perkins, M.W., Govindarajulu, P., Seglie, D., Walker, S., Cunningham, A.A. & Fisher, M.C. (2006). The emerging amphibian pathogen *Batrachochytrium dendrobatidis* globally infects introduced populations of the North American bullfrog, *Rana catesbeiana*. *Biology Letters* 2, 455–459.
- Garner, T.W.J., Garcia, G., Carroll, B. & Fisher, M.C. (2009b). Using itraconazole to clear *Batrachochytrium dendrobatidis* infection, and subsequent depigmentation of *Alytes muletensis* tadpoles. *Diseases of Aquatic Organisms* 83, 257–260.
- Garner, T.W.J., Walker, S., Bosch, J., Leech, S., Rowcliffe, J.M., Cunningham, A.A. & Fisher, M.C. (2009a). Life history tradeoffs influence mortality associated with the amphibian pathogen *Batrachochytrium dendrobatidis*. *Oikos* 118, 783–791.
- Garner, T.W.J., Martel, A., Bielby, J., Bosch, J., Anderson, L.G., Meredith, A., Cunningham, A.A., Fisher M.C., Henk, D.A. & Pasmans, F. (2013). Infectious diseases that may threaten Europe's amphibians. In: *Amphibian Biology Volume 11: Status of Conservation and Decline of Amphibians: Eastern hemisphere, Part 3: Western Europe*. Eds: Heatwole, H. & Wilkinson, J.W., Pelagic Publishing, Exeter, United Kingdom.
- Gibbons, J.W., Scott, D.E., Ryan, T.J., Buhlmann, K.A., Tuberville, T.D., Metts, B.S., Greene, J.L., Mills, T., Leiden, Y., Poppy, S. & Winne, C.T. (2000). The global decline of reptiles, Déjà Vu Amphibians: reptile species are declining on a global scale. Six significant threats to reptile populations are habitat loss and degradation, introduced invasive species, environmental pollution, disease, unsustainable use, and global climate change. *AIBS Bulletin* 50, 653–666.
- González, D. L., Baláž, V., Solský, M., Thumsová, B., Kolenda,

- K., Najbar, A., Kautman, M., Chajma, R., Balogová, M. & Vojar, J. (2019). Recent findings of potentially lethal salamander fungus *Batrachochytrium salamandrivorans*. *Emerging Infectious Diseases* 25(7), 1416-1418.
- Grasselli, E., Bianchi, G., Dondero, L., Marchianò, V., Carafa, M., Perrone, M. & Salvidio, S. (2019). First screening for *Batrachochytrium salamandrivorans* (Bsal) in wild and captive salamanders from Italy. *Salamandra* 5(2), 124-126.
- Gray, M.J., Duffus, A.L.J., Haman, K.H., Harris, R.N., Allender, M.C., Thompson, T.A., Christman, M.R., Sacerdote-Velat, A., Sprague, L.A., Williams, J.M. & Miller, D.L. (2017). Pathogen Surveillance in Herpetofaunal Populations: Guidance on Study Design, Sample Collection, Biosecurity, and Intervention Strategies. *Herpetological Review* 48, 334-351.
- Grogan, L.F., Robert, J., Berger, L., Skerratt, L.F., Scheele, B.C., Castley, J.G., Newell, D.A. & McCallum, H. I. (2018). Review of the Amphibian Immune Response to Chytridiomycosis, and Future Directions. *Frontiers in Immunology* 9, 2536.
- Hidalgo-Vila, J., Díaz-Paniagua, C., Marchand, M.A. & Cunningham, A.A. (2012). *Batrachochytrium dendrobatidis* infection of amphibians in the Doñana National Park, Spain. *Diseases of Aquatic Organisms* 98, 113-119.
- Holopainen, R., Ohlemeyer, S., Schütze, H., Bergmann, S.M. & Tapiovaara, H. (2009). Ranavirus phylogeny and differentiation based on major capsid protein, DNA polymerase and neurofilament triplet H1-like protein genes. *Diseases of Aquatic Organisms* 85, 81-91.
- Hyatt, A.D., Gould, A.R., Zupanovic, Z., Cunningham, A.A., Henstberger, S., Whittington, R.J., Kattenbelt, J. & Coupar, B.E.H. (2000). Comparative studies of piscine and amphibian iridoviruses. *Archives of Virology* 145, 301-331.
- Jakob-Hoff R.M., MacDiarmid S.C., Lees C., Miller P.S., Travis D. & Kock R. (2014). – *Manual of Procedures for Wildlife Disease Risk Analysis*. World Organisation for Animal Health, Paris, 160 pp. Published in association with the International Union for Conservation of Nature and the Species Survival Commission.
- Kärvemo, S., Meurling, S., Berger, D., Höglund, J. & Laurila, A. (2018). Effects of host species and environmental factors on the prevalence of *Batrachochytrium dendrobatidis* in northern Europe. *PLoS ONE* 13, e0199852.
- Kik, M., Martel, A., Spitzen-van der Sluijs, A., Pasmans, F., Wohlsein, P., Gröne, A. & Rijks, J.M. (2011). Ranavirus-associated mass mortality in wild amphibians, The Netherlands, 2010: A first report. *The Veterinary Journal* 190, 284-286.
- Kolenda, K., Najbar, A., Ogielska, M. & Balá, V. (2017). *Batrachochytrium dendrobatidis* is present in Poland and associated with reduced fitness in wild populations of *Pelodytes lessonae*. *Diseases of Aquatic Organisms* 124, 241-245.
- Laking, A.E., Ngo, H.N., Pasmans, F., Martel, A. & Nguyen, T.T. (2017). *Batrachochytrium salamandrivorans* is the predominant chytrid fungus in Vietnamese salamanders. *Scientific Reports* 7, 44443
- Langwig, K.E., Voyles, J., Wilber, M.Q., Frick, W.F., Murray, K.A., Bolker, B.M., Collins, J.P., Cheng, T.L., Fisher, M.C., Hoyt, J.R. et al. (2015). Context-dependent conservation responses to emerging wildlife diseases. *Frontiers in Ecology and the Environment* 13, 195-202.
- Lorch, J. M., Lankton, J., Werner, K., Falendysz, E. A., McCurley, K. & Blehert, D. S. (2015). Experimental infection of snakes with *Ophidiomyces ophidiicola* causes pathological changes that typify snake fungal disease. *mBio* 6, e01534-15.
- Lorch, J. M., Knowles, S., Lankton, J. S., Michell, K., Edwards, J.L., Kapfer, J. M., Staffen, R. A., Wild, E.R., Schmidt, K.Z., Ballmann, A. E., & Blodgett, D. (2016). Snake fungal disease: an emerging threat to wild snakes. *Philosophical Transactions of the Royal Society B* 371, e20150457.
- Madsen, T., Stille, B. & Shine, R. (1996). Inbreeding depression in an isolated population of adders *Vipera berus*. *Biological Conservation* 75, 113-118.
- Mali, I., Villamizar-Gomez, A., Krizmanić, I., Ajtić, R. & Forstner, M.R. (2017). Evidence of *Batrachochytrium dendrobatidis* Infection in Amphibians from Serbian Lowlands. *Journal of Wildlife Diseases* 53, 686-689.
- Marantelli, G., Berger, L., Speare, R. & Keegan, L. (2004). Distribution of the amphibian chytrid *Batrachochytrium dendrobatidis* and keratin during tadpole development. *Pacific Conservation Biology* 10, 173-179.
- Marschang, R.E. (2011). Viruses infecting reptiles. *Viruses* 3, 2087-2126.
- Martel, A., Fard, M.S., Van Rooij, P., Jooris, R., Boone, F., Haesebrouck, F., Van Rooij, D. & Pasmans, F. (2012). Road-killed common toads (*Bufo bufo*) in Flanders (Belgium) reveal low prevalence of ranaviruses and *Batrachochytrium dendrobatidis*. *Journal of Wildlife Diseases* 48, 835-839.
- Martel, A., Spitzen-van der Sluijs, A., Blooi, M., Bert, W., Ducatelle, R., Fisher, M.C., Woeltjes, A., Bosman, W., Chiers, K., Bossuyt, F. & Pasmans, F. (2013). *Batrachochytrium salamandrivorans* sp. nov. causes lethal chytridiomycosis in amphibians. *Proceedings of the National Academy of Sciences*, 201307356.
- Martel, A., Blooi, M., Adriaensen, C., Van Rooij, P., Beukema, W., Fisher, M.C., Farrer, R.A., Schmidt, B.R., Tobler, U., Goka, K., Lips, K.R., Muletz, C., Zamudio, K.R., Bosch, J. Lötters, S., Wombwell, E., Garner, T.W. J., Cunningham, A.A., Spitzen-van der Sluijs, A., Salvidio, S., Ducatelle, R., Nishikawa, K., Nguyen, T.T., Kolby, J.E., Van Bocxlaer, I., Bossuyt, F. & Pasmans, F. (2014). Recent introduction of a chytrid fungus endangers Western Palearctic salamanders. *Science* 346, 630-631.
- Martínez-Solano, I., Bosch, J., & García-París, M. (2003). Demographic trends and community stability in a montane amphibian assemblage. *Conservation Biology* 17, 238-244.
- Miaud, C., Pozet, F., Gaudin, N.C.G., Martel, A., Pasmans, F. & Labrut, S. (2016). *Ranavirus* causes mass die-offs of alpine amphibians in the southwestern Alps, France. *Journal of Wildlife Diseases* 52, 242-252.
- Meier, G., Notomista, T., Marini, D. & Ferri, V. (2018). First case of Snake Fungal Disease affecting a free-ranging *Natrix natrix* (Linnaeus, 1758) in Ticino Canton, Switzerland. *Herpetology Notes* 11, 885-891.
- Mutchmann, F. & Scheenweiss, D. (2008). Herpes virus infektionen bei *Pelodytes fuscus* und anderen anuren in Berlin-Brandenburg Raum. *Rana Sonderheft* 5, 113-118.
- Nazir, J., Spengler, M. & Marschang, R. E., 2012. Environmental persistence of amphibian and reptilian ranaviruses. *Diseases of Aquatic Organisms* 98, 177-184.
- Nguyen, T.T., Van Nguyen, T., Ziegler, T., Pasmans, F. & Martel, A. (2017). Trade in wild anurans vectors the urodelan pathogen *Batrachochytrium salamandrivorans* into Europe.

- Amphibia-Reptilia* 38, 554-556.
- Nichols, D. K., Lamirande, E. W., Pessier, A. P. & Longcore, J. E. (2001). Experimental transmission of cutaneous chytridiomycosis in dendrobatid frogs. *Journal of Wildlife Diseases* 37, 1-11.
- O'Hanlon, S. J., Rieux, A., Farrer, R.A., Rosa, G.M., Waldman, B., Bataille, A., Kosch, T.A., Murray, K.A., Brankovics, B., Fumagalli, M., Martin, M.D., Wales, N., Alvarado-Rybak, M., Bates, K.A., Berger, L. et al. (2018). Recent Asian origin of chytrid fungi causing global amphibian declines. *Science* 360, 621-627.
- Ohst, T., Gräser, Y., Mutschmann, F. & Plötner, J. (2011). Neue Erkenntnisse zur Gefährdung europäischer Amphibien durch den Hautpilz *Batrachochytrium dendrobatidis*. *Zeitschrift für Feldherpetologie* 18, 1-17.
- Ohst, T., Gräser, Y. & Plötner, J. (2013). *Batrachochytrium dendrobatidis* in Germany: distribution, prevalences, and prediction of high risk areas. *Diseases of Aquatic Organisms* 107, 49-59.
- Olson, D.H., Aanensen, D.M., Ronnenberg, K.L., Powell, C.I., Walker, S.F., Bielby, J., Garner, T.W.J., Weaver, G. & Fisher, M.C. (2013). Mapping the global emergence of *Batrachochytrium dendrobatidis*, the amphibian chytrid fungus. *PLoS ONE* 8, e56802.
- Origi, F.C., Schmit, B.R., Lohman, P., Otten, P., Akdesir, A., Gashen, V., Aguilar-Bultet, L., Wahli, T., Sattler, U. & Stoffel, M.H. (2017). *Ranid herpesvirus 3* and proliferative dermatitis in free-ranging wild common frogs (*Rana temporaria*). *Veterinary Pathology* 54, 686-694.
- Origi, F.C., Schmit, B.R., Lohman, P., Otten, P., Meier, R.K., Pisano, S.R.R., Moore-Jones, G., Tecilla, M., Sattler, U., Wahli, T., Gashen, V. & Stoffel, M.H. (2018). *Bufo* *herpesvirus 1* (BfHV1) associated dermatitis and mortality in free ranging common toads (*Bufo bufo*) in Switzerland. *Scientific Reports* 8, 14737.
- Ouellet, M., Dejean, T. & Galois, P. (2012). Occurrence of the amphibian chytrid fungus *Batrachochytrium dendrobatidis* in introduced and native species from two regions of France. *Amphibia-Reptilia* 33, 415-422.
- Paré, J. A. & Sigler, L. (2016). An overview of reptile fungal pathogens in the genera *Nannizziopsis*, *Paranannizziopsis*, and *Ophidiomyces*. *Journal of Herpetological Medicine and Surgery* 26, 46-53.
- Pasmans, F., Mutschmann, F., Halliday, T. & Zwart, F. (2006). Amphibian declines: the urgent need for amphibian research in Europe. *Veterinary Journal* 171, 18-19.
- Pasmans, F., Muijsers, M., Maes, S., Van Rooij, P., Brutyn, M., Ducatelle, R., Haesebrouck, F. & Martel, A. (2010). Chytridiomycosis related mortality in a midwife toad (*Alytes obstetricans*) in Belgium. *Vlaams Diergeneeskundig Tijdschrift* 79, 460-462.
- Pasmans, F., Van Rooij, P., Blooi, M., Tessa, G., Bogaerts, S., Sotgiu, G., Trenton Garner, T.W.J., Fisher, M.C., Schmidt, B.R., Woeltjes, T. et al. (2013). Resistance to chytridiomycosis in European plethodontid salamanders of the genus *Speleomantes*. *PLoS ONE* 8, e63639.
- Patrelle, C., Miaud, C., Cristina, N., Kulberg, P. & Merilä, J. (2012). Chytrid Fungus Screening in a Population of Common Frogs from Northern Finland. *Herpetological Review* 43, 422-425
- Price, S.J., Garner, T.W.J., Nichols, R.A., Balloux, F., Ayres, C., de Alba, A.M.C. & Bosch, J. (2014). Collapse of amphibian communities due to an introduced *Ranavirus*. *Current Biology* 24, 2586-2591.
- Price, S.J., Wadia, A., Wright, O.N., Leung, W.T., Cunningham, A.A. & Lawson, B. (2017). Screening of a long-term sample set reveals two *Ranavirus* lineages in British herpetofauna. *PLoS ONE* 12, e0184768.
- Rajeev, S., Sutton, D.A., Wickes, B.L., Miller, D.L., Giri, D., Van Meter, M., Thompson, E.H., Rinaldi, M.G., Romanelli, A.M., Cano, J.F. & Guarro, J. (2009). Isolation and characterization of a new fungal species, *Chrysosporium ophidiicola*, from a mycotic granuloma of a black rat snake (*Elaphe obsoleta obsoleta*). *Journal of Clinical Microbiology* 47, 1264-1268.
- Rijks, J.M., Saucedo, B., Spitzen-van der Sluijs, A., Wilkie, G.S., van Asten, A.J., van den Broek, J., Boonyarittichaij, R., Stege, M., van der Sterren, F., Martel, A. & Pasmans, F. (2016). Investigation of amphibian mortality events in wildlife reveals an on-going ranavirus epidemic in the north of the Netherlands. *PLoS One* 11(6), p.e0157473.
- Rosa, G.M., Anza, I., Moreira, P.L., Conde, J., Martins, F., Fisher, M.C. & Bosch, J. (2013). Evidence of chytrid-mediated population declines in common midwife toad in Serra da Estrela, Portugal. *Animal Conservation* 16, 306-315.
- Rosa, G.M., Sabino-Pinto, J., Laurentino, T.G., Martel, A., Pasmans, F., Rebelo, R., Griffiths, R.A., Stöhr, A.C., Marschang, R.E., Price, S.J. & Garner, T.W.J. (2017). Impact of asynchronous emergence of two lethal pathogens on amphibian assemblages. *Scientific Reports* 7, 43260.
- Sabino-Pinto, J., Bletz, M., Hendrix, R., Perl, R.B., Martel, A., Pasmans, F., Lötters, S., Mutschmann, F., Schmeller, D.S., Schmidt, B.R. et al. (2015). First detection of the emerging fungal pathogen *Batrachochytrium salamandrivorans* in Germany. *Amphibia-Reptilia* 36, 411-416.
- Sabino-Pinto, J., Krause, E.T., Bletz, M.C., Martel, A., Pasmans, F., Steinfartz, S. & Vences, M. (2018). Detectability vs. time and costs in pooled DNA extraction of cutaneous swabs: a study on the amphibian chytrid fungi. *Amphibia-Reptilia* 40, 29-39 <https://doi.org/10.1163/15685381-20181011>.
- Saucedo, B., Hughes, J., Spitzen-Van Der Sluijs, A., Kruithof, N., Schills, M., Rijks, J.M., Jacinto-Maldonado, M., Suarez, N., Haenen, O.L., Voorbergen-Laarman, M. & Van Den Broek, J. (2018). *Ranavirus* genotypes in the Netherlands and their potential association with virulence in water frogs (*Pelophylax* spp.). *Emerging Microbes & Infections* 7, 1-14.
- Saucedo, B., Garner, T.W.J., Kruithof, N., Allain, S.J.R., Goodman, M.J., Cranfield, R.J., Sergeant, C., Vergara, D.A., Kik, M.J.L., Forzán, M.J., van Beurden, S.J. & Gröne, A. (2019). Common midwife toad ranaviruses replicate first in the oral cavity of smooth newts (*Lissotriton vulgaris*) and show distinct strain-associated pathogenicity. *Scientific Reports* 9, 4453.
- Scalera, R., Adams, M.J. & Galvan, S.K. (2008). Occurrence of *Batrachochytrium dendrobatidis* in amphibian populations in Denmark. *Herpetological Review* 39, 199-200.
- Scheele, B.C., Pasmans, F., Skerratt, L.F., Berger, L., Martel, A., Beukema, W., Acevedo, A.A., Burrowes, P.A., Carvalho, T., Catenazzi, A. et al. (2019). Amphibian fungal panzootic causes catastrophic and ongoing loss of biodiversity. *Science* 363:1459-1463.
- Scott, M.E. (1988). The impact of infection and disease on animal populations: implications for conservation biology. *Conservation Biology* 2, 40-56.
- Sharifian-Fard, M., Pasmans, F., Adriaenssens, C., Devisscher, S.,

- Adriaens, T., Louette, G. & Martel, A., (2011). Ranaviriosis in invasive bullfrogs, Belgium. *Emerging Infectious Diseases* 17, 2371-2372.
- Sigler, L., Hambleton, S. & Paré, J. A. (2013). Molecular characterization of reptile pathogens currently known as members of the *Chysosporium* anamorph of *Nannizziopsis vriesii* (CANV) complex and relationship with some human-associated isolates. *Journal of Clinical Microbiology* 51, 3338-3357.
- Simoncelli, F., Fagotti, A., Dall'Olio, R., Vagnetti, D., Pascolini, R. & Di Rosa, I. (2005). Evidence of *Batrachochytrium dendrobatidis* infection in water frogs of the *Rana esculenta* complex in central Italy. *EcoHealth* 2, 307.
- Skerratt, L.F., Berger, L., Speare, R., Cashins, S., McDonald, K.R., Phillott, A.D., Hines, H.B. & Kenyon, N. (2007). Spread of chytridiomycosis has caused the rapid global decline and extinction of frogs. *EcoHealth* 4, 125-134.
- Sodhi, N.S., Bickford, D., Diesmos, A.C., Lee, T.M., Koh, L.P., Brook, B.W., Sekercioglu, C.H. & Bradshaw, C.J. (2008). Measuring the meltdown: drivers of global amphibian extinction and decline. *PLoS one* 3, e1636.
- Spitzen-van der Sluijs, A., Spikmans, F., Bosman, W., de Zeeuw, M., van der Meij, T., Goverse, E., Kik, M., Pasmans, F. & Martel, A. (2013). Rapid enigmatic decline drives the fire salamander (*Salamandra salamandra*) to the edge of extinction in the Netherlands. *Amphibia-Reptilia* 34, 233-239.
- Spitzen-van der Sluijs, A., Martel, A., Hallmann, C.A., Bosman, W., Garner, T. W. J., Van Rooij, P., Jooris, R., Haesebrouck, F. & Pasmans, F. (2014). Environmental determinants of recent endemism of *Batrachochytrium dendrobatidis* infections in amphibian assemblages in the absence of disease outbreaks. *Conservation Biology* 28, 1302-1311.
- Spitzen-van der Sluijs, A., Martel, A., Asselberghs, J., Bales, E.K., Beukema, W., Bletz, M.C., Dalbeck, L., Goverse, E., Kerres, A., Kinet, T. & Kirst, K. (2016a). Expanding distribution of lethal amphibian fungus *Batrachochytrium salamandrivorans* in Europe. *Emerging Infectious Diseases* 22, 1286.
- Spitzen-van der Sluijs, A., van den Broek, J., Kik, M., Martel, A., Janse, J., van Asten, F., Pasmans, F., Gröne, A. & Rijks, J.M. (2016b). Monitoring Ranavirus-associated mortality in a Dutch heathland in the aftermath of a *Ranavirus* disease outbreak. *Journal of Wildlife Diseases* 52, 817-827.
- Spitzen-van der Sluijs, A., Pasmans, F., Struijk, R.P., Schils, M., Doornbos, P., van der Sterren, F., Rijks, J., Kik, M., Saucedo, B., Bosman, W. & Martel, A. (2016c). Course of an isolated *Ranavirus* outbreak in a *Pelobates fuscus* population in the Netherlands. *Journal of Herpetological Medicine and Surgery* 26, 117-121.
- Spitzen-van der Sluijs, A., Canessa, S., Martel, A. & Pasmans, F. (2017). Fragile coexistence of a global chytrid pathogen with amphibian populations is mediated by environment and demography. *Proceedings of the Royal Society B: Biological Sciences* 284, 20171444.
- Spitzen-van der Sluijs, A., Stegen, G., Bogaerts, S., Canessa, S., Steinfartz, S., Janssen, N., Bosman, W., Pasmans, F. & Martel, A. (2018). Post-epizootic salamander persistence in a disease-free refugium suggests poor dispersal ability of *Batrachochytrium salamandrivorans*. *Scientific Reports* 8, 3800.
- Stagni, G., Dall'olio, R., Fusini, U., Mazzotti, S., Scocciati, C. & Serra, A. (2004) Declining populations of Apennine yellow-bellied toad *Bombina pachypus* in the northern Apennines (Italy): is *Batrachochytrium dendrobatidis* the main cause? *Italian Journal of Zoology* 71, 151-154.
- Stegen, G., Pasmans, F., Schmidt, B.R., Rouffaer, L.O., Van Praet, S., Schaub, M., Canessa, S., Laudelout, A., Kinet, T., Adriaensen, C., Haesebrouck, F., Bert, W., Bossuyt, F. & Martel, A. (2017). Drivers of salamander extirpation mediated by *Batrachochytrium salamandrivorans*. *Nature* 544, 353-356.
- Stöhr, A.C., Hoffmann, A., Papp, T., Robert, N., Pruvost, N.B., Reyer, H.U. & Marschang, R.E. (2013). Long-term study of an infection with ranaviruses in a group of edible frogs (*Pelophylax kl. esculentus*) and partial characterization of two viruses based on four genomic regions. *The Veterinary Journal* 197, 238-244.
- Stöhr, A.C., López-Bueno, A., Blahak, S., Caeiro, M.F., Rosa, G.M., de Matos, A.P.A., Martel, A., Alejo, A. & Marschang, R.E. (2015). Phylogeny and differentiation of reptilian and amphibian ranaviruses detected in Europe. *PLoS ONE* 10, e0118633.
- Stuart, S.N., Chanson, J.S., Cox, N.A., Young, B.E., Rodrigues, A. S., Fischman, D.L. & Waller, R.W. (2004). Status and trends of amphibian declines and extinctions worldwide. *Science* 306, 1783-1786.
- Sura, P., Janulis, E. & Profus, P. (2010). Chytridiomikoza – śmiertelne zagrożenie dla płazów. *Chrońmy Przyrodę Oczyszczą* 66, 406-421.
- Sztatecsny, M. & Glaser, F. (2011). From the eastern lowlands to the western mountains: first records of the chytrid fungus *Batrachochytrium dendrobatidis* in wild amphibian populations from Austria. *The Herpetological Journal* 21, 87-90.
- Tan, W.G., Barkman, T.J., Chinchar, V.G. & Essani, K., (2004). Comparative genomic analyses of frog virus 3, type species of the genus *Ranavirus* (family Iridoviridae). *Virology* 323, 70-84.
- Teacher, A.G.F., Cunningham, A.A. & Garner, T.W.J. (2010). Assessing the long-term impact of *Ranavirus* infection in wild common frog populations. *Animal Conservation* 13(5), 514-522.
- Thomas, V., Blooi, M., Van Rooij, P., Van Praet, S., Verbrugghe, E., Grasselli, E., Lukac, M., Smith, S., Pasmans, F., and Martel, A., (2018). Recommendations on diagnostic tools for *Batrachochytrium salamandrivorans*. *Transboundary and Emerging Diseases* 65, e478-e488.
- Tinsley, R.C., Coxhead, P.G., Stott, L.C., Tinsley, M.C., Piccinni, M.Z. & Guille, M.J. (2015). Chytrid fungus infections in laboratory and introduced *Xenopus laevis* populations: assessing the risks for UK native amphibians. *Biological Conservation* 184, 380-388.
- Tobler, U. & Schmidt, B. R. (2010). Within-and among-population variation in chytridiomycosis-induced mortality in the toad *Alytes obstetricans*. *PLoS One* 5, e10927.
- Tobler, U., Borgula, A. & Schmidt, B.R. (2012). Populations of a susceptible amphibian species can grow despite the presence of a pathogenic chytrid fungus. *PLoS ONE* 7, e34667.
- Todd, B.D., Willson, J.D. & Gibbons, J.W. (2010). The global status of reptiles and causes of their decline. In: *Ecotoxicology of Amphibians and Reptiles*, 2nd Eds: Sparling, D.W., Bishop,

- C.A., & Krest, S., CRC Press, Boca Raton, USA.
- Tünde, G.J., Szabó, K. & Vörös, J. (2012). Kitridiomikózis vizsgálata egy magas-bakonyi vizes élőhely kétéltűközösségén. *Állattani Közlemények* 97, 47-59.
- Ursenbacher, S., Monney, J.C. & Fumagalli, L. (2009). Limited genetic diversity and high differentiation among the remnant adder (*Vipera berus*) populations in the Swiss and French Jura Mountains. *Conservation Genetics* 10, 303-315.
- van Beurden, S.J., Hughes, J., Saucedo, B., Rijks, J., Kik, M., Haenen, O.L., Engelsma, M.Y., Gröne, A., Verheije, M.H., & Wilkie, G. (2014). Complete genome sequence of a common midwife toad virus-like *Ranavirus* associated with mass mortalities in wild amphibians in the Netherlands. *Genome Announcements* 2, e01293-14.
- Van Rooij, P., Martel, A., Haesebrouck, F., & Pasmans, F. (2015). Amphibian chytridiomycosis: a review with focus on fungus-host interactions. *Veterinary Research* 46, 137.
- Van Rooij, P., Pasmans, F., Coen, Y. & Martel, A. (2017). Efficacy of chemical disinfectants for the containment of the salamander chytrid fungus *Batrachochytrium salamandrivorans*. *PLoS ONE* 12, e0186269.
- Vojar, J., Havlíková, B., Solský, M., Jablonski, D., Iković, V. & Baláž, V. (2017). Distribution, prevalence and amphibian hosts of *Batrachochytrium dendrobatidis* in the Balkans. *Salamandra* 53, 44-49.
- Vörös, J., Bosch, J., Dán, Á. & Hartel, T. (2013). First record of *Batrachochytrium dendrobatidis* on amphibians in Romania. *North-Western Journal of Zoology* 9, 446-449.
- Wake, D.B. & Vredenburg, V.T. (2008). Are we in the midst of the sixth mass extinction? A view from the world of amphibians. *Proceedings of the National Academy of Sciences* 105, 11466-11473.
- Weldon, C., Du Preez, L.H., Hyatt, A.D., Muller, R. & Speare, R. (2004). Origin of the amphibian chytrid fungus. *Emerging Infectious Diseases* 10, 2100-2105.
- Wood, L.R., Griffiths, R.A. & Schley, L. (2009). Amphibian chytridiomycosis in Luxembourg. *Bulletin de la Société des Naturalistes Luxembourgeois* 110, 109-114.
- Woodhams, D.C., Bigler, L. & Marschang, R. (2012). Tolerance of fungal infection in European water frogs exposed to *Batrachochytrium dendrobatidis* after experimental reduction of innate immune defenses. *BMC Veterinary Research* 8, 197.
- World Organisation for Animal Health (OIE) & International Union for Conservation of Nature (IUCN) (2014). – Guidelines for Wildlife Disease Risk Analysis. OIE, Paris, 24 pp. Published in association with the IUCN and the Species Survival Commission.
- Zampiglia, M., Canestrelli, D., Chiocchio, A. & Nascetti, G. (2013). Geographic distribution of the chytrid pathogen *Batrachochytrium dendrobatidis* among mountain amphibians along the Italian peninsula. *Diseases of Aquatic Organisms* 107, 61-68.

Accepted: 3 April 2019